# Strategies for Organic and Low-input Integrated Breeding and Management

## Newsletter 1

#### Table of contents

October 2011

• WP1. Identify traits for organic/low-input systems p.2 •WP2. Identify and monitor genetic diversity p.3 •WP3. Exploitation of diversity in breeding p.5 •WP4. Exploitation of diversity in management p.6 •WP5. Comparison of conventional/organic breeding p.8 •WP6. Participatory plant breeding and management p.9 • WP7. Breeding/management effects on crop quality p.11 • WP8. Sustainability assessment of innovations p.12 • WP9. Dissemination, training and technology transfer p.13 • Deliverables and calendar p.14 • SOLIBAM snapshots p.15 • SOLIBAM partners p.16

> Project Coordinator Veronique Chable (INRA) veronique.chable@rennes.inra.fr

In charge of dissemination activities Riccardo Bocci (AIAB) r.bocci@aiab.it

**Collaborative Project** (large-scale integrating project)



The SOLIBAM project is supported by the **European Commission through the Seventh** Framework Programme for Research and Technological Development under the Grant Agreement n°245058



# The rationale of SOLIBAM

research

Veronique Chable (INRA - Project co-ordinator)

This is the first in a series of newsletters, which aim to outline the activities of the SOLIBAM consortium. The overall objective of SOLIBAM is to develop specific and novel breeding approaches integrated with management practices to improve the performance, quality, sustainability and stability of crops adapted to organic and low-input

systems. The project i s founded on the concept of diversity. Our research

hypothesis is that

developing diversity at all levels is the best strategy for improving the capacity for crops to adapt to fluctuating environmental conditions and for increasing yields and yield stability in organic and low-input systems. Diversity is also a characteristic of the consortium members, who bring a wide range of competencies and scientific approaches, farming systems experience, climatic conditions and cultural backgrounds. The partners come from 12 countries, representing many different regions from the North of Europe to Africa and our consortium represents most of

the disciplines needed to explore and sustain diversity in the field.

During this first stage of the project, we have succeeded in building a cohesive team approach. The diversity of members has led to the development of what we may call: "the SOLIBAM strategy". From the first annual meeting, the definition of our



strategy incorporated both scientific n Ь а organisational viewpoints: In SOLIBAM we are

combining many disciplines and values with the aim of increasing system diversity and participatory methods. In addition to developing techniques and assessing quantitative and qualitative data, transdisciplinary thinking is central to the success of SOLIBAM.

In this first newsletter, we offer an overview of all the disciplines involved in the development of an innovative sustainable strategy for producing quality crops and appropriate processes for the food supply system based on PPBM (Participatory Plant Breeding and Management) and diversity at all levels.

### Identify traits for organic/low-input systems

Workpackage 1

#### Frederic Rey (ITAB)

## Diversity of contexts is fundamental in Solibam: a national chapter for each country involved

Describing the range of geographical and cultural backgrounds to organic and low input crop production is one goal of the Solibam project, as this kind of data has not been collected yet, neither in Europe or in other Solibam member countries. The national chapters provide a first step to defining the various situations in terms of agro-ecological conditions, seed regulation and the organic seed sector. All partners have contributed to this activity by completing a "common questionnaire" in 2010. The compilation of this data is in process and is being carried out by ITAB, the lead organisation for this task (T1.1).

#### A survey of stakeholder's expectation

There is a lack of varieties adapted to organic systems in the agricultural sector. Within the framework of Solibam (T1.1), a survey was undertaken to determine the important traits specific to organic and low input farming for cereals including wheat, and for horticultural crops including tomatoes. The survey also aims to describe organic stakeholders' expectations in plant breeding over a wide range of different agro-climatic conditions in Europe and Africa. The

result of this study will be to provide data where existing published information is scarce, for example: identification of the varieties currently grown by the organic/LI producers; reasons for their variety choice; producers' expectations for organic varieties. Most of the published papers dealing with organic plant breeding programs concern cereal crops (Lammerts



et al. 2008, Goldringer et al. 2010). Papers on horticultural crops are limited and those on organic stakeholders expectations are even fewer. The recent State of Organic Seed report (USA) is an exception which can nevertheless be highlighted; based on stakeholder surveys it analyses the challenges and opportunities in building the organic seed sector (Dillon & Hubbard 2011).

In our study, as a first step, a qualitative survey (with open questions) was implemented in six different countries: Ethiopia, France, Germany, Italy, Mali and Portugal. This work aimed to identify the key research questions/hypotheses within both a European and African perspective as useful plant traits for organic/LI agriculture may differ considerably from one country to another depending on the agro-ecological conditions and on the market. As an example, two main market categories have been clearly identified from the French stakeholders' answers. These categories and results fit with the hypothesis that the market is a significant factor influencing the choice of seeds and varieties. Expectations and practices of producers selling on a local market (i.e. direct sale) differ radically

> from those of producers selling to large retailers. For vegetables, the impact of nurseries is considered as a major factor, and this issueneeds more investigations. The final analysis of this work is expected in late 2012.

> > Tomato seedlings tested in Italy in WP3

Acknowledgments: The following persons are involved in this WP1 (T1.1) work; they have contributed to the methodology and have implemented this survey in their own countries: Riccardo Bocci (Italy, Associazione Italiana per l'Agricoltura Biologica), Hans-Jürgen Reents (Germany, Technical University of Munich), Isabel Dinis, Daniela Santos and Pedro Mendes Moreira (Portugal, Escola Superior Agraria de Coimbra), Salif Foulani Sissoko (Mali, Coordination Nationale des Organisations Paysannes du Mali), Fetien Abay and Abdelkadir Kedir (Ethiopia, Mekele University).

#### References

Dillon M, Hubbard K (2011): State of organic seed. Organic seed alliance report http://www.seedalliance.org/Publications/, (accessed 2011-06-01).

Goldringer I., Dawson J., Vettoretti A., Rey F. (2010): Breeding for resilience: a strategy for organic and low-input farming systems?. Eucarpia 2<sup>nd</sup> conference of the "Organic and Low-Input" Section, 1-3 Dec. 2010, Paris, France <u>http://orgprints.org/18171/1/Breeding for resilience%2DBook of abstracts.pdf</u>, (accessed 2011-06-01).

Lammerts E., Ostegard H., Goldringer I., Scholten O. (2008): Plant breeding for organic and sustainable, low input agriculture: dealing with genotype-environment interactions, Euphytica, Vol. 163, Number 3, 546 p.

Lammerts van Bueren ET, Struik PC, Jacobsen E (2002): Ecological concepts in organic farming and their consequences for an organic crop ideotype. Netherlands Journal of Agricultural Science 50 (1): 1-26

Wolfe MS, Baresel JP, Desclaux D, Goldringer I, Hoad S, Kovacs G, Löschenberger F, Miedaner T, Østergård H, Lammerts Van Bueren ET. (2008): Developments in breeding cereals for organic agriculture in Europe. Euphytica 163:323–346



#### Identify and monitor genetic diversity Workpackage 2

#### Isabelle Goldringer (INRA)

farming implies working with complex traits (QTL mapping), WP2 focuses on populations and controlled by many genes interacting with each plants developed for organic or LI systems under other and with complex and variable specific systems (i.e. plant breeding scheme x crop environments, new approaches are needed to management) in WP3, 5 & 6. appropriately integrate the most recent biotechnology tools into specific breeding for organic and LI conditions. WP2 objective is to renew the genetic concepts of plant breeding for organic and LI agriculture, by associating knowledge from genomics, quantitative genetics, population genetics and epigenetic approaches to phenotyping under different organic and lowinput crop management systems. In contrast to the usual approaches where associations between molecular markers and phenotypes are studied in

Because breeding for organic or low-input (LI) a specific design optimized for their detection

In Task 2.1, WP2 is identifying and checking for polymorphisms in breeding populations developed in WP3, WP5 & WP6 using three relevant types of molecular markers already available: (i) neutral or background DNA markers, (ii) markers related to epigenetic states and, (iii) markers in candidate genes. Different species important in the SOLIBAM project are being studied: wheat (winter and durum wheat), barley, maize, bean, broccoli and, tomato. Individuals from maize, bean, barley, wheat and broccoli populations have been



Illumina GoldenGate Output: Multiple SNPs at Once (G. Pea, SSSUP)

variability of outcrossing in populations will be assessed in relation to the different agroecological conditions in order to develop strategies to increase and maintain heterozygosity a n d heterogeneity, and thus the buffering capacity, within the

sampled and DNA extraction is crop management systems x completed or in progress. agro-climatic conditions to Depending on the species, better understand the responses microsatellite or SNP will be of different kind of populations used as background neutral (e.g. composite cross markers. Illumina Golden Gate populations vs mixtures) to assay has been tested by SSSUP different selection practices, and partner on the parents of a CCP to develop strategies for populations (WP3) and it gave a maintaining appropriate levels of change. rate of diversity within the populations/ polymorphism. The list of key varieties. Their efficiency in candidate genes for the different monitoring the evolution of species is under development diversity under selection will be based on the literature. Markers evaluated and compared. of the epigenetic state Original association methods

(INRA).

satisfactory

In Task 2.2, the resulting polymorphic markers will be used to assess the evolution of genetic diversity under different



Wheat seedlings - INRA Le Moulon (France)

(methylation) are under will be developed to apply development in broccoli within and between the (UNIPG) and in wheat for a breeding populations in order to specific vernalization gene identify genetic and epigenetic markers associated with phenotypic responses to selection pressures (agroclimatic conditions, crop management systems, selection

> practices). Phenotypic responses will be assessed considering both specific adaptive traits (such as drought resistance) and more general adaptation features. The potential role of a few selected candidate genes will be further analysed through transcriptional characterisation of the genes in contrasting genotypes. The levels and

populations, increasing the potential for movement of plants between agroecological zones in response to climate

In Task 2.3, multivariate approaches to selection responses and methods for association studies in breeding populations will be developed. Based on the insights obtained on the genetic and epigenetic mechanisms involved in adaptation to organic and LI farming, appropriate strategies incorporating marker assisted selection (MAS) at different levels will be proposed. Multivariate approaches will be used to assess the selection strength of various traits under different conditions and to predict selection responses. The results obtained on the molecular diversity of all populations will be discussed during a common meeting at month M40 with all the partners that have been involved in the development of the populations in other WPs (e.g. WP3 and WP5) and also including the partners who are not directly involved in WP2.

### Exploitation of diversity in breeding Workpackage 3

#### Sally Howlett (ORC)

The focus of Solibam WP3 is the 'Exploitation of Diversity in Breeding'. This work package incorporates trials of a number of vegetable and arable crops in a wide range of countries. The objective is to increase and utilise diversity in crops in both organic and low-input systems in order to improve their yield and quality at the same time as dealing with increasing environmental variability. Diversity encompasses

both novel sources, such as synthetically created lines with new genomic structures, and existing sources, such as landraces, variety mixtures and populations; all are included in the Solibam approach. The relative strengths and weaknesses in performance of crops containing differing levels of diversity are being compared in the WP3 trials. The findings will also feed into other workpackages, including those centred on elucidating the underlying genetics of adaptation (WP2), the role of management in utilising diversity (WP4) and evaluating the organoleptic characteristics of the crops produced (WP7).

A key driver for WP3 is the need for crops bred specifically for organic and low-input systems.Currently, commercially available varieties have largely been developed for performance

under high-input conditions and they often fail to achieve acceptable and reliable yields in more heterogeneous organic and low-input environments. Furthermore, increasingly unstable weather patterns put more demands on crops, which require a greater capacity for adaptability if they are to be resilient to and perform reliably under such conditions.

The crops involved are broccoli, common bean, cabbage, tomatoes, maize and cereals (winter wheat, durum wheat, einkorn, emmer and barley). The trials include a large number of Solibam partners in three subtasks:

#### A) Population development

Novel populations are being developed in broccoli (calabrese type), durum, einkorn, emmer and bread wheat using different techniques to intercross a wide range of genetic resources. The populations will be trialled against modern lines; this part of the WP is led by Véronique Chable (INRA) and Geza Kovacs (HAS). Broccoli trials are already underway at three French sites and the

Wheat - evolutionary population (Italy)

novel cereal populations are currently being bred and multiplied for inclusion in project years 2-4 (2011-12).

B) Comparison of breeding strategies

A larger number of crops are included in comparisons of breeding strategies in winter wheat, barley, common bean, sprouting broccoli, cabbage, maize and tomato. Led by ORC, the trials compare genotypes representing different breeding strategies (populations, mixtures, landraces and pedigree lines) in varied geographical locations in Europe and sub-Saharan Africa. A major assumption to be tested is that reliability will be correlated with increased within-crop diversity. Trials started in 2010 and will continue through to project year 4.

C) Integrating the new and existing approaches

The idea here is to integrate the best populations with the best in pedigree line breeding. The trials focus on winter wheat and are being carried out in Hungary (HAS), Austria (DONAU) and the UK (ORC) in all project years. The hope is that mixing pure line varieties with populations will simultaneously combine the benefits of high performance and diversity in a flexible way, which could also extend the useful life of a variety

#### Exploiting diversity in crop management Workpackage 4

#### Paolo Bàrberi (SSSUP)

diversification within and between crop species, management and habitat, coupled with use and development of genetically diverse germplasm. performance analysis of highly diverse germplasm different pedo-climatic conditions in Europe and the sustainability analysis (WP8). Sub-Saharan Africa. The degree of diversification and typology of interventions differ from place to place depending on the prioritisation of crop features as well as on local opportunities and constraints, but common treatments across sites are being used wherever possible.

WP4 is structured into four tasks. The first aims to design novel organic and low-input cropping systems based on optimum use of agrobiodiversity by simultaneously taking into account the genotype, environment and management components altogether. The hypothesis is that cropping systems based on higher genotype, species and/or management diversity would increase crop yield, yield stability, produce quality, soil fertility and weed suppression. Experiments will make use of genetically diverse germplasm (wheat, maize and tomato) developed by SOLIBAM partners. When



SSSUP wheat trials - Pisa (Italy)

agrobiodiversity at the species and/or variety level and management diversity on arbuscular-

Our goal is to design, develop and test innovative sustainability of the systems. A common set of arable and vegetable cropping systems based on a parameters will be measured across all experiments high level of agrobiodiversity. This means to ensure homogeneous evaluation of cropping system effects on key traits. Data and samples collected in WP4 experiments will feed the global Our activities are deployed across a range of (WP1), the analysis of produce quality (WP7) and



SSSUP wheat trials - Pisa (Italy)

The second task aims to improve performance of diversified cropping systems through (co)breeding. Plant breeding has always considered selection under monoculture conditions, thus little is known about populations grown in competition. We expect that there will be a strong selection pressure towards better adaptation to these conditions. The objective of this task is to identify genotypes adapted to diversified systems and to assess changes in morphophysiological and genetic characteristics of populations. We are identifying traits that contribute to higher competitive ability, which are necessary to obtain higher yield under living mulch systems, and we will select the best performing genotypes in cereal-legume systems. Work is being carried out on wheat, maize and common bean.

highly diverse germplasm is not available, increased The third task is checking the effects of breeding (e.g. crop rotation, cover crops, intercropping, mycorrhizal fungi (AMF). The research hypothesis variety mixtures) and at the management/habitat is that cropping systems with higher breeding and/ level (e.g. tillage, sowing method, weed or management diversity are more prone to management, non cropped areas) will compensate develop effective AMF symbiosis, thereby for this. These experiments will highlight the contributing to soil fertility enhancement in feasibility of using cropping system diversification organic and low-input agriculture. We will identify as an effective strategy to improve crop how certain cultivars of model species of cereals performance, quality traits, and the overall and grain legumes have a better ability to associate



with AMF and form effective symbiosis, and how plant breeding methodologies may take this trait into account. Also, we are testing if diversified cropping systems including legumes as cover crops or intercrops increase AMF root colonisation of host crops and associated agroecosystem services, and whether AMF inoculation can be used as a management strategy to enhance crop productivity.

The final task is testing the effect of conservation and/or (re)introduction, and of appropriate management of linear and non linear seminatural elements (e.g. hedgerows, field margins, small woodland patches) on production-related agroecosystem services (biological control of pests and prevention of weed invasion). The research hypothesis is that the maintenance of well structured non cultivated areas (i.e. functional biodiversity at the habitat level) promote production-related agroecosystem services and that this effect is greater in more diversified cropped fields. It is expected that the results will also provide hints on which types of structure and management of non cropped areas (e.g. flora composition, habitat complexity, mowing and/or cultivation of field margins) are more likely to enhance these agroecosystem services.



Old wheat varieties trials - France

Comparison of conventional/organic breeding Workpackage 5

#### Geza Kovàcs (HAS)

The focus of Solibam WP5 is to compare the initiated under different environments and methods and the efficiency of organic breeding approaches with conventional breeding methods to be able to improve the competitiveness of organic varieties in the most appropriate way. The work package involves characterisation of the most important genetic resources of a number of autogamous and allogamous species for their possible use in organic breeding, followed by studies of the effectiveness of early stage selection under different management systems. The data obtained will support WP1 activities and selected materials will be incorporated to the different experiments and trials within WP3 and WP4. During the study of the effectiveness of early stage selection several new crosses have been developed in a large number of species, and the selection work was

management conditions in order to study their effects and interactions. The performance and yield stability of the existing varieties of bread wheat and durum wheat grown under organic and conventional systems was studied using ring tests in three different countries, and detailed studies on the resistance against common bunt were carried out. According to the time schedule the majority of the experiments are at the initial phase of development, but the first selection steps have been successfully completed. Based on the results from these existing varieties, the development of organic maintenance breeding methods has been initiated, and tests of the effect of post harvest methods for organic seed production were carried out to examine the whole process of organic variety production.

Farm days in Italy in June 2001. In the left Castellaneta (Puglia), in the right Isernia (Molise).





#### Participatory plant breeding and management Workpackage 6

#### Salvatore Ceccarelli and Adnan Al Yassin (ICARDA)

The objective of WP6 is to develop participatory plant breeding and management strategies to address the specific demands from marginal areas or small scale agriculture in developing countries, Europe and Africa. The difference between participatory research applied to plant breeding and management (PPBM), and conventional research is that in PPBM scientists offer farmers, end-users and consumers a number of choices (varieties, populations, management options, combinations of management options and varieties) so that farmers can select, after an adequate period of testing under their own conditions, the most appropriate option. Therefore, PPBM takes into account the diversified management practices, needs, expectations and traditions of farmers' preferences and is particularly suitable for the development of new varieties and management practices adapted to low input and organic farming. One innovative aspect that WP6 has introduced is evolutionary participatory plant breeding which is based on populations with large genetic variability, including mixtures of landraces, which can be handled directly by farmers in a multitude of environments while they slowly evolve and adapt. It is like making available, in the hands of farmers, an evolving gene bank of a given crop. One major advantage of evolutionary breeding is its simplicity and its

enormous potential to adapt crops – any crop – to climatic changes as well as other agronomic changes which might occur in the future (Ceccarelli et al., 2010). The expected outputs of WP6 are improved germplasm, populations and/ or varieties, specifically adapted to these conditions that will be more easily adopted by farmers.

WP6 is organised into three components:

In Task 6.1 we analysed important experiences in participatory research. This component has assembled an inventory of 22 PPBM cases covering Wheat (5), Legumes (3), Cauliflower, Barley (6), Maize (3), Tomato, Faba bean (2), Sweet Potato, Cassava, Sorghum (3), Rice (1) Millet (1), Fonio (1) and Potato from the UK, France (2), Ethiopia, Portugal, Spain (2), Uganda, Ghana, Burkina Faso, Mali, Algeria, Iran, Eritrea, Nepal (2), Yemen, Netherlands, Nicaragua and China. The inventory has been assembled mostly through informal contacts and is currently going through a critical analysis by the partners. The results of this analysis will be useful to implement activities in other work packages.

In Task 6.2 we develop methodologies for participatory research: one of the activities has been the preparation of a Manual on PPB which will help to disseminate the methodologies for this approach. Training courses based on the manual have been conducted in a number of

Farm days in Italy in June 2001. On the left Caltagirone (Sicilia), On the right Peccioli (Toscana).





countries, including Ethiopia, with The participation of University of Mekelle. The manual covers a summary of the main breeding methodologies of self-, cross-, and vegetativelypropagated crops. It describes both participatory plant breeding and participatory variety selection, and discusses the detailed description of possible experimental designs and statistical analysis. The final section of the manual presents examples of multi-environment data analysis, genotype x environment interactions, variety release and seed production and examples of PPB impacts. These methodologies have been implemented in trials planted in 16 locations in Ethiopia in 2010/2011. The trials were designed at ICARDA and three types of germplasm were planted across 12 farmers' fields, of which 5 were conducted in women's farms due to their direct involvement in the decision making process of the varietal selection, and 6 Farmers Training centres. Three evolutionary populations, one on barley (based on about 1600 F2), one on durum wheat (based on

### On the left: Farmers discussing barley attributes during field selection day

Bottom: The participants to the training in data analysis: from left to right Mr. Mulken Bantayehu, Mr. Dejene Kassahun, Mr. G/ Egzihabher Miruts, Mr. Molla Fentie, Mr. Afework Kebede, Dr. Ceccarelli, Dr. Berhane Lakew, Mr. Yosef Gebrehawariyat. Sitting Mr. Tamene Mideksa and Mr. Kefyalew Taye.



about 700 F2) and one on bread wheat (based on more than 2000 segregating populations) have been planted in many locations in Italy (Toscana, Molise, Puglia and Sicilia). Sufficient amounts of seed were harvested to distribute the populations to other farmers practising different management systems.

The objectives of Task 6.3 are (i) to identify the effects of farmer mass selection in several species and in different conditions and (ii) to measure the evolution and the adaptation of the populations in terms of genetic and phenotypic diversity. During the first period, field experiments have been set up for most species (wheat, barley, maize, bean and broccoli) in France, Italy, Portugal, Ethiopia and Mali. Germplasm has been distributed to farmers, measures have been recorded and trials have been harvested. An experiment to assess farmers' mass selection response has been carried out in France on bread wheat and the first results indicate that farmers' mass selection within populations has a significant positive impact on traits of interest. This will be assessed for more populations, other species and countries. The impact of the participatory breeding programme implementation was assessed as very positive in Ethiopia.



### Breeding/management effects on crop quality Workpackage 7

#### Marianna Rakszegi (HAS)

Organoleptic, nutritional and end-use properties of the breeding lines and populations developed under WP3-6 will be studied in the framework of WP7 taking into consideration the global and local needs and expectations of the consumers. Properties of plants growing at different agro-climatic conditions will be studied using different breeding techniques and crop management systems. Hedonic tests with a large panel of consumers as well as sensory analysis with trained juries will be carried out in order to select the organic cereal, legume and vegetable genotypes preferred by consumers. Nutritional properties, such as antioxidant levels, protein, starch and amylose content will also be measured to evaluate the health related effects of the organically bred lines. At the same time the impact of crop management and breeding approach (crop rotation, mulching, irrigation etc.) on the end-use quality will be studied (e.g. bread-making, pasta making), particularly taking into consideration the requirements for the production of special local products. Results will ultimately support the breeders, the agronomists, the processing industry and last, but not least, the consumers.

In the first period of the project, a methodology was developed to integrate organoleptic quality criteria in breeding programs of tomatoes, broccoli and bread. This resulted in a tasting guide which recommends four different tests and explains how to prepare samples and analyse the results. This guide gives a common starting point for all those partners who are planning to study organoleptic quality of vegetables or bread.

Adaptation and optimisation of extraction and analytical methods have also been started. Aroma and flavor compounds will be measured by SPME-GC-MS, while carotenoids, tocopherols and phenolic compounds will be analysed by HPLC.

Barley genotypes were selected for kolo production in Ethiopia and their nutritional quality ( $\beta$ -glucan, Fe, Zn) was studied. One genotype with outstanding iron (Fe) content was identified. Laboratories are awaiting the arrival of seeds expected from the harvest of 2011

#### Sustainability assessment of innovations Workpackage 8

Michal Kulak (FDEA-ART), Mads Ville Markussen and Hanne Østergård (Risø-DTU), Laurence Smith (ORC) and Thomas Nemecek (FDEA-ART)

Ιn work package 8 we are assessing the sustainability of a range of farm management strategies using 8 case study farms. In order to do this, we are evaluating current practices, and the resultant environmental and socio-economic impacts from the implementation of SOLIBAM strategies. We also want to know what farmers views are of the SOLIBAM strategies in



addition to the food supply and legislationrelated issues that can influence their adoption. The methods that are being used include Life Cycle Assessment, Energy and eMergy analysis and Cost Benefit Analysis.

The assessments are based on a number of farm and food supply case studies located in Europe, selected according to pre-defined criteria. The samples represent various European climatic conditions, different production systems and marketing schemes. What is common to all cases is a willingness to reduce farm external inputs as much as possible and to produce food in a sustainable manner.

With help from local partners, the team established contact with the case-study farms. During the visits, data on external inputs, production, financial data and distribution channels have been recorded. So far we have been to France to look at grain and bread production and in UK to investigate vegetable production and distribution.

The first French case is a small 5 ha farm that was established only 10 years ago. The main product for the farm is bread, produced using grain that is grown and milled on the farm.

The other farm is a well established farm of more than 70 ha that, in addition to its cereals and flour production, also produces milk and a variety of cheeses. Both farms are selling directly to end consumers and both are involved in the participatory breeding of cereals.

In the UK we have visited two stockless vegetable producers; a large farm that supplies most of its products to pack houses that distribute the vegetables to supermarkets and a small market garden with more than 70 different (!) vegetables on its 7 hectares.

With the first datasets gathered, the next step will be to feed the data into our different models while the data collection continues on more case study farms throughout Europe.

One of the biggest challenges is to adapt existing models for sustainability assessment to these highly diverse farms. As for other work packages within SOLIBAM it is a challenge to conduct research that is both relevant to the project and society in general, and is multidisciplinary in its approach.

### Dissemination, training and technology transfer Workpackage 9

#### Riccardo Bocci (AIAB)

SOLIBAM dissemination activities in the first year have been focused primarily on setting up the website, designing and publishing the brochure in 6 languages and organising the first round of Farm days. Partners have also participated in many international and national meetings in order to present the project and its objectives.

Farm days (see table 1 for 2010 activities) are designed to be our main dissemination tool among farmer's communities in each country. They will be held each year, to enable



breeders, farmers, extension services and researchers involved in WPs to share their skills, information, and knowledge with participating and non-participating farmers. In addition, Farm days act as a forum for the discussion of project results and related topics with farmers. ITAB is in charge of the overall co-ordination of Farm days.

Table 1 - 2010 Farm days in the different countries					
Country	Date 2010	Visit	Species	People involved	
		Research Regional platform on farmers' varieties (Association AgroBio Périgord)		Farmers, association's animators, researchers, local politicians	
		Salon Ille et Bio ( by the association Culture Bio and the Triptolème)	Wheat	Farmers, association's animators and volunteers, researchers, local politicians	
	10th	Fête de la vache nantaise (involvement of the association Triptolème and INRA in the fair)		Farmers, researchers, association animators and volunteers	
Austria	June 30th	Visit to the farm of Gerhard Hof		Saatbau Linz ( <u>www.saatbaulinz.at</u> ), seed producing farmers for Saatbau Linz, and Saatzucht Donau	
Portugal	June 4th	APCC association		Some farmers involved in the trials, the farmer (Mr. Francisco Meireles who still run the PPB programme since 1984) and the Breeder Dr. Silas Pego who started the VASO PPB programme in 1984	
UK	June 15th	Wakelyns Agroforestry Open Day	Beans	Local farmers and growers, ORC staff.	
	June 1st	Visit to the farm of Rosario Floriddia (collection of wheat landraces).	Wheat, barley	Farmers, Public institutions, researchers	
Italy	June 2nd	Meeting at the farm of Oriana Porfiri (collection of wheat, barley and emmer landraces).		Farmers, Public institutions, researchers	
	June 4th		Radicchio (red-leafed chicory, barley, wheat	Farmers, Public institutions, researchers	

SOLIBAM has also started activities regarding training and dissemination in Africa. ICARDA and the University of Mekelle conducted a training course on participatory plant breeding in Ethiopia (April 2010). Twelve scientists from Universities and National and Regional Agricultural Centres attended the training course.

### List of deliverables published

We provide a complete list of the deliverables released until now by the SOLIBAM consortium. The ones marked with PU are public and freely downloadable from our website <u>www.solibam.eu</u>.



Download the brochure

- D1.1 Methodologies to assess farmer-perceived needs and practices (PU)
- D1.5 Determination of traits most relevant for model species (PU)
- D9.1 Public WebSite (PU)
- D9.2 Project brochure (PU)
- D9 4 Printing of the Tasting Guide booklet (PU)
- D9.7 Report on the outcomes of the farm days in 2010 (PU)
- D10.1 KOM minutes
- D10.2 collaborative workspace
- D10.3 3rd Executive Committee Meeting minutes
- D10.3 2nd Executive Committee Meeting minutes
- D10.4 1st Annual meeting minutes



### Next meetings and events

15 -16 September - Pisa (IT) - SOLIBAM WP4 Workshop
26-27 September - Seoul (South Korea) - IFOAM Organic seed pre-conference
28 September - 1 October - Seoul (South Korea) - IFOAM Organic World Congress
18-19 October - Wakelyns (UK) - SOLIBAM Executive committee
3-4 November - Frankfurt (DE) - European Consortium for Organic Plant Breeding 10 year's Anniversary Conference. Organic Plant Breeding: What makes the difference?
20-22 November - Rennes (FR) - SOLIBAM WP8 Workshop



# The 23 SOLIBAM Partners

Institut National de la Recherche Agronomique (France)	(IIII) INRA				
Associazione Italiana per l'Agricoltura Biologica (Italy)					
The Organic Research Centre, Elm Farm (UK)	OKCANE EXTARCH				
RISØ National Laboratory for Sustainable Energy (Denmark)					
Institut Technique de l'Agriculture Biologique (France)	TAB				
Technical University of Munich (Germany)	TUT				
Instituto de Tecnologia Quimica e Biologica (Portugal)	mob par				
Agencia Estatal Consejo Superior de Investigaciones Cientificas/Instituto de Agricultura Sostenible <mark>(Spain)</mark>	CSIC				
Escola Superior Agraria de Coimbra (Portugal)					
Agricultural Research Institute of the Hungarian Academy of Sciences (Hungary)	NU				
Scuola Superiore Sant'Anna, Pisa (Italy)	Scuola Superiore Sant'Anna				
University of Perugia (Italy)					
Agroscope Reckenholz-Taenikon Research Station ART (Switzerland)	Consistent and the second seco				
Institute of Food and Resource Economics (Denmark)					
INRA Transfert (France)	INRA Transfert				
University of Pisa (Italy)	UNIVERSITA' DI PISA				
Crop breeding companies					
Saatzucht Donau - cereal breeding (Austria)					
Gautier Semence - vegetable breeding (France)	GAUTIER				
Agrovegetal - legume breeding (Spain)	where the				
Arcoiris - vegetable breeding (Italy)	And the second				
Institutions from African countries and international organisation					
International Center for Agricultural Research in the Dry Areas (International)	4				
Coordination Nationale des Organisations Paysannes du Mali (Mali)					
Mekelle University (Ethiopia)					



http://www.solibam.eu

For information: <u>r.bocci@aiab.it</u>